

Nouvelles perspectives sur l'énumération des classes de permutations

Michael Albert

Department of Computer Science, University of Otago

Sep 7, 2012, LaBRI



New perspectives on the enumeration of permutation classes

Michael Albert

Department of Computer Science, University of Otago


Sep 7, 2012, LaBRI



Permutation classes


Definition

A **permutation class** is a collection of permutations, \mathcal{C} , with the property that, if $\pi \in \mathcal{C}$ and we erase some points from its plot, then the permutation defined by the remaining points is also in \mathcal{C} .



492713685 $\in \mathcal{C}$

implies



21543 $\in \mathcal{C}$



Permutation classes

- ▶ Permutation classes are simply downward closed sets in the *subpermutation* ordering



Permutation classes

- ▶ Permutation classes are simply downward closed sets in the *subpermutation* ordering
- ▶ The objective is to try to *understand* the structure of permutation classes (or to identify when this is possible)



Permutation classes

- ▶ Permutation classes are simply downward closed sets in the *subpermutation* ordering
- ▶ The objective is to try to *understand* the structure of permutation classes (or to identify when this is possible)
- ▶ Enumeration is a consequence or symptom of such understanding



Permutation classes

- ▶ Permutation classes are simply downward closed sets in the *subpermutation* ordering
- ▶ The objective is to try to *understand* the structure of permutation classes (or to identify when this is possible)
- ▶ Enumeration is a consequence or symptom of such understanding
- ▶ If X is a set of permutations, then $\text{Av}(X)$ is the permutation class consisting of those permutations which do not dominate any permutation of X



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $\text{Av}(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)
- ▶ $\text{Av}(1234)$ was enumerated by Gessel (1990)



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)
- ▶ $Av(1234)$ was enumerated by Gessel (1990)
- ▶ $Av(1342)$ by Bóna (1997)



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)
- ▶ $Av(1234)$ was enumerated by Gessel (1990)
- ▶ $Av(1342)$ by Bóna (1997)
- ▶ The enumeration of $Av(1324)$ is still unknown



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)
- ▶ $Av(1234)$ was enumerated by Gessel (1990)
- ▶ $Av(1342)$ by Bóna (1997)
- ▶ The enumeration of $Av(1324)$ is still unknown
- ▶ For single element bases of length 5 or more, nothing much is known in the non-monotone case



First steps

- ▶ It seems plausible that the fewer permutations there are in a class, or alternatively the more restrictive the conditions of membership, then the more likely it is to have good structure
- ▶ Simion and Schmidt (1985) successfully enumerated all classes $Av(X)$ for $X \subseteq \mathcal{S}_3$
- ▶ For a single avoided pattern of length 4, it turns out there are only three possible enumerations (based on both specific and general symmetries)
- ▶ $Av(1234)$ was enumerated by Gessel (1990)
- ▶ $Av(1342)$ by Bóna (1997)
- ▶ The enumeration of $Av(1324)$ is still unknown
- ▶ For single element bases of length 5 or more, nothing much is known in the non-monotone case
- ▶ All doubleton bases of lengths 4 and 3, and many of length 4 and 4 are known



Stanley-Wilf conjecture

Relative to the set of all permutations, proper permutation classes are small. Specifically:

Theorem

Let \mathcal{C} be a proper permutation class. Then, the growth rate of \mathcal{C} ,

$$\text{gr}(\mathcal{C}) = \limsup |\mathcal{C} \cap \mathcal{S}_n|^{1/n}$$

is finite.

This was known as the *Stanley-Wilf conjecture* and it was proven in 2004 by Marcus and Tardos.


The obvious next questions are:

- ▶ What growth rates can occur?
- ▶ What can be said about classes of particular growth rates?



Antichains

The subpermutation order contains infinite antichains.



Consequently, there exist 2^{\aleph_0} distinct enumeration sequences for permutation classes – we must be careful not to try to do too much.



Small growth rates

- ▶ Kaiser and Klazar (EJC, 2003) showed that the only possible values of $\text{gr}(\mathcal{C})$ less than 2 are the greatest positive solutions of:

$$x^k - x^{k-1} - x^{k-2} - \dots - x - 1 = 0$$



Small growth rates

- ▶ Kaiser and Klazar (EJC, 2003) showed that the only possible values of $\text{gr}(\mathcal{C})$ less than 2 are the greatest positive solutions of:

$$x^k - x^{k-1} - x^{k-2} - \dots - x - 1 = 0$$

- ▶ Vatter (Mathematika, 2010) showed that every real number larger than the unique real solution (approx. 2.482) of

$$x^5 - 2x^4 - 2x^2 - 2x - 1 = 0$$

occurs as a growth rate



Small growth rates

- ▶ Kaiser and Klazar (EJC, 2003) showed that the only possible values of $\text{gr}(\mathcal{C})$ less than 2 are the greatest positive solutions of:

$$x^k - x^{k-1} - x^{k-2} - \dots - x - 1 = 0$$

- ▶ Vatter (Mathematika, 2010) showed that every real number larger than the unique real solution (approx. 2.482) of

$$x^5 - 2x^4 - 2x^2 - 2x - 1 = 0$$

occurs as a growth rate

- ▶ V (PLMS, 2011) further showed that the smallest growth rate of a class containing an infinite antichain is the unique positive solution, $\kappa \simeq 2.20557$ of

$$x^3 - 2x^2 - 1 = 0$$

and completely characterized the set of possible growth rates below κ



Simple permutations

Definition

A permutation is **simple** if it contains no nontrivial consecutive subsequence whose values are also consecutive (though not necessarily in order)




Simple permutations

Definition

A permutation is **simple** if it contains no nontrivial consecutive subsequence whose values are also consecutive (though not necessarily in order)

2 5 7 4 6 1 3



Not simple




Simple permutations

Definition

A permutation is **simple** if it contains no nontrivial consecutive subsequence whose values are also consecutive (though not necessarily in order)

2 5 7 3 6 1 4



Simple



Simple permutations

Definition

A permutation is **simple** if it contains no nontrivial consecutive subsequence whose values are also consecutive (though not necessarily in order)

- ▶ Simple permutations form a positive proportion of all permutations (asymptotically $1/e^2$)
- ▶ In many (conjecturally all) proper permutation classes they have density 0
- ▶ We can hope to understand a class by understanding its simples and how they *inflate*
- ▶ Specifically, this may yield functional equations of the generating function and hence computations of the enumeration and/or growth rate



Finitely many simple permutations

Theorem

If a class has only finitely many simple permutations then it has an algebraic generating function.

- ▶ A and Atkinson (2005)
- ▶ Effective ‘in principle’, i.e. an algorithm for computing a defining system of equations for the generating function
- ▶ Some interesting corollaries, e.g. if a class has finitely many simples and does not contain arbitrarily long decreasing permutations then it has a rational generating function
- ▶ *“The prime reason for giving this example is to show that we are not necessarily stymied if the number of simple permutations is infinite.”*



Encodings

- ▶ The enumerative combinatorics of words and trees is better understood



Encodings

- ▶ The enumerative combinatorics of words and trees is better understood
- ▶ So, apply leverage from there by encoding permutation classes



Encodings

- ▶ The enumerative combinatorics of words and trees is better understood
- ▶ So, apply leverage from there by encoding permutation classes
- ▶ For trees this approach goes by many names



Encodings

- ▶ The enumerative combinatorics of words and trees is better understood
- ▶ So, apply leverage from there by encoding permutation classes
- ▶ For trees this approach goes by many names
- ▶ For words, A, At and Ruškuc (2003) give (fairly) general criteria for the encoding of a permutation class by a regular language





Encodings

- ▶ The enumerative combinatorics of words and trees is better understood
- ▶ So, apply leverage from there by encoding permutation classes
- ▶ For trees this approach goes by many names
- ▶ For words, A, At and Ruškuc (2003) give (fairly) general criteria for the encoding of a permutation class by a regular language
- ▶ Extended to a new type of encoding, the *insertion encoding* (prefigured by Viennot) by A, Linton and R (2005)



Grid classes

- ▶ The notion of *griddable* class was central to V's characterization of small permutation classes
- ▶ Loosely, a griddable class is associated with a matrix whose entries are (simpler) permutation classes
- ▶ All permutations in the class can be chopped apart into sections that correspond to the matrix entries




Geometric monotone grid classes

In a *geometric* grid class, the permutations need to be drawn from the points of a particular representation in \mathbb{R}^2

Theorem (A, At, Bouvel, R and V (to appear TAMS))

Every geometrically griddable class:

- ▶ *is partially well ordered;*
- ▶ *is finitely based;*
- ▶ *is in bijection with a regular language and thus has a rational generating function.*



Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):

Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):

- ▶ Let $\mathcal{C}[\mathcal{D}]$ represent the inflations of permutations in \mathcal{C} by permutations in \mathcal{D} , and let $\langle \mathcal{C} \rangle$ denote the closure of \mathcal{C} under inflation

Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):

- ▶ Let $\mathcal{C}[\mathcal{D}]$ represent the inflations of permutations in \mathcal{C} by permutations in \mathcal{D} , and let $\langle \mathcal{C} \rangle$ denote the closure of \mathcal{C} under inflation
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ is finitely based and partially well ordered

Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):

- ▶ Let $\mathcal{C}[\mathcal{D}]$ represent the inflations of permutations in \mathcal{C} by permutations in \mathcal{D} , and let $\langle \mathcal{C} \rangle$ denote the closure of \mathcal{C} under inflation
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ is finitely based and partially well ordered
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ has an algebraic generating function

Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):

- ▶ Let $\mathcal{C}[\mathcal{D}]$ represent the inflations of permutations in \mathcal{C} by permutations in \mathcal{D} , and let $\langle \mathcal{C} \rangle$ denote the closure of \mathcal{C} under inflation
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ is finitely based and partially well ordered
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ has an algebraic generating function
- ▶ If \mathcal{C} is geometrically griddable and \mathcal{U} is strongly rational, then $\mathcal{C}[\mathcal{U}]$ is also strongly rational

Beyond grid classes

Results from *Inflations of Geometric Grid Classes of Permutations*, A, R and V (arxiv.org/abs/1202.1833):


- ▶ Let $\mathcal{C}[\mathcal{D}]$ represent the inflations of permutations in \mathcal{C} by permutations in \mathcal{D} , and let $\langle \mathcal{C} \rangle$ denote the closure of \mathcal{C} under inflation
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ is finitely based and partially well ordered
- ▶ If \mathcal{C} is geometrically griddable, then every subclass of $\langle \mathcal{C} \rangle$ has an algebraic generating function
- ▶ If \mathcal{C} is geometrically griddable and \mathcal{U} is strongly rational, then $\mathcal{C}[\mathcal{U}]$ is also strongly rational
- ▶ Every small permutation class has a rational generating function

Quasi-applications

- ▶ These ideas, together with a certain amount of number eight wire or duct tape, i.e. “un peu de rafistolage” can be used to compute enumerations for some (arguably) interesting classes
- ▶ Examples from the basic environment of geometric grid classes are considered in A, At and Brignall: *The Enumeration of Three Pattern Classes using Monotone Grid Classes* (EJC 19.3 (2012) P20)
- ▶ Examples for inflations of geometric grid classes are considered in A, At and V: *Inflations of Geometric Grid Classes: Three Case Studies* (arxiv.org/abs/1209.0425)

Av(4312, 3142)

- ▶ Every simple permutation in this class lies in the geometric grid class:



- ▶ This yields a regular language for the simple permutations
- ▶ The allowed inflations of these permutations are easily described, yielding a recursive description of the class
- ▶ This leads to an equation for its generating function:

$$\begin{aligned}(x^3 - 2x^2 + x)f^4 &+ (4x^3 - 9x^2 + 6x - 1)f^3 \\ &+ (6x^3 - 12x^2 + 7x - 1)f^2 \\ &+ (4x^3 - 5x^2 + x)f \\ &+ x^3 &= 0\end{aligned}$$

Av(321)

This class is in some sense a limit of geometric grid classes:




Av(321)

This class is in some sense a limit of geometric grid classes:




$Av(321)$

This class is in some sense a limit of geometric grid classes:




Av(321)

This class is in some sense a limit of geometric grid classes:




Av(321)

This class is in some sense a limit of geometric grid classes:




Av(321)

This class is in some sense a limit of geometric grid classes:




Av(321)

This class is in some sense a limit of geometric grid classes:



$Av(321)$

This class is in some sense a limit of geometric grid classes:



Conjecture

Every finitely based proper subclass of $Av(321)$ has a rational generating function

Where to from here?

- ▶ Underlying the main results on geometric grid classes and their inflations is a notion of *natural encoding*, in this case of permutations by words, which may be applicable to other types of combinatorial structure particularly those carrying a linear order

Where to from here?

- ▶ Underlying the main results on geometric grid classes and their inflations is a notion of *natural encoding*, in this case of permutations by words, which may be applicable to other types of combinatorial structure particularly those carrying a linear order
- ▶ Understanding the limit case for geometric grid classes better

Where to from here?

- ▶ Underlying the main results on geometric grid classes and their inflations is a notion of *natural encoding*, in this case of permutations by words, which may be applicable to other types of combinatorial structure particularly those carrying a linear order
- ▶ Understanding the limit case for geometric grid classes better
- ▶ Understanding non-geometric grid classes (David Bevan, a student of R. Brignall has some nice results in the case where there is only one cycle, and is working on more general cases)

Where to from here?

- ▶ Underlying the main results on geometric grid classes and their inflations is a notion of *natural encoding*, in this case of permutations by words, which may be applicable to other types of combinatorial structure particularly those carrying a linear order
- ▶ Understanding the limit case for geometric grid classes better
- ▶ Understanding non-geometric grid classes (David Bevan, a student of R. Brignall has some nice results in the case where there is only one cycle, and is working on more general cases)
- ▶ *Permutation Patterns 2013*: July 1-5, Paris