


The ascent lattice on Dyck paths

with Jean-Luc Baril, Sergey Kirgizov
(LIB, Dijon) and Mehdi Naima (LIP6, Paris)





arXiv:2409.15982

Mireille Bousquet-Mélou
CNRS, LaBRI, Université de Bordeaux

Outline


- Orders on Dyck paths --- The ascent order
- Lattice properties (the Nadeau-Tewari lattice)
- Counting intervals
- m -Dyck paths, and mirrored m -Dyck paths
- Connection with the Sylvester congruence [Hivert, Novelli, Thibon 05]

I. Orders on Dyck paths





Dyck paths

- A Dyck path of **size** $n=8$ (size=number of up steps)



Dyck paths


- A Dyck path of **size** $n=8$ (size=number of up steps)



Stanley's lattice S_n

A poset on Dyck paths of size n

Def. The path u is smaller than v if u lies below v .




Stanley's lattice S_n

A poset on Dyck paths of size n

Def. The path u is smaller than v if u lies below v .

- Cover relations are given by $EN \rightarrow NE$ (that is, valley \rightarrow peak)




Stanley's lattice S_n

A poset on Dyck paths of size n

Def. The path u is smaller than v if u lies below v .

- Cover relations are given by $EN \rightarrow NE$ (that is, valley \rightarrow peak)
- A sub-lattice of the Young lattice




Stanley's lattice S_n


A poset on Dyck paths of size n

Def. The path u is smaller than v if u lies below v .

- Cover relations are given by $EN \rightarrow NE$ (that is, valley \rightarrow peak)
- A sub-lattice of the Young lattice
- Graded (by area)




- A poset on Dyck paths with n up steps
- Cover relations: choose a valley in the path u .
Swap the East step and the **shortest Dyck path** that follows.



The Tamari lattice

[Tamari 5]

- A poset on Dyck paths with n up steps
- Cover relations: choose a valley in the path u .
Swap the East step and the **shortest Dyck path** that follows.




The Tamari lattice

[Tamari 5]

- A poset on Dyck paths with n up steps
- Cover relations: choose a valley in the path u .

Swap the East step and the **shortest Dyck path** that follows.




The Tamari lattice

[Tamari 5]


- A poset on Dyck paths with n up steps
- Cover relations: choose a valley in the path u .

Swap the East step and the **shortest Dyck path** that follows.




The ascent poset (or: greedy Stanley?)

- A poset on Dyck paths with n up steps
- Cover relations: choose a valley in the path u . Swap the down step and the **ascent** that follows. (the path moves up)




[Chenevière, Nadeau...]


Ascent posets: $n = 3, 4$



Ascent posets: $n = 3, 4$




Poset inclusion



A characterization of the ascent order

Proposition. In the ascent poset, $P \preceq Q$ iff


- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



A characterization of the ascent order

Proposition. In the ascent poset, $P \preceq Q$ iff


- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



A characterization of the ascent order

Proposition. In the ascent poset, $P \preceq Q$ iff

- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



Applications:

- ◆ lattice structure
- ◆ recursive construction of intervals

II. Lattice structure

(Each pair of elements has a sup and an inf)

The Nadeau-Tewari poset


[2024]

Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if

- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .

$$v = (10, 10, 10, 10, 9, 4)$$

$$u = (10, 10, 8, 8, 7, 3)$$





The Nadeau-Tewari poset

[2024]

Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if

- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .





The Nadeau-Tewari poset

[2024]

Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if

- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .




The Nadeau-Tewari poset

[2024]


Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if

- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .



$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$

$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$




The Nadeau-Tewari poset

[2024]

Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if


- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .

Observation: Dyck paths of size n can be encoded by nonincreasing sequences of length n , and then the ascent order coincides with the NT order.




$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$

$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$



The Nadeau-Tewari poset and the ascent order

Observation: The ascent order is the order induced by the NT order on sequences $u=(u_1, \dots, u_n)$ such that $n-i+1 \leq u_i \leq n$ for all i .




$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$

$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$

The Nadeau-Tewari poset and the ascent order

Observation: The ascent order is the order induced by the NT order on sequences $u=(u_1, \dots, u_n)$ such that $n-i+1 \leq u_i \leq n$ for all i .

These sequences form an **interval** with $u_{\min} = (n, n-1, \dots, 1)$ and $u_{\max} = (n, n, \dots, n)$.



$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$


$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$

The Nadeau-Tewari poset and the ascent order

Observation: The ascent order is the order induced by the NT order on sequences $u=(u_1, \dots, u_n)$ such that $n-i+1 \leq u_i \leq n$ for all i .

These sequences form an **interval** with $u_{\min} = (n, n-1, \dots, 1)$ and $u_{\max} = (n, n, \dots, n)$.

Proposition [NT 24]: the NT poset is a **lattice**.



$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$

$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$


The Nadeau-Tewari poset and the ascent order

Observation: The ascent order is the order induced by the NT order on sequences $u=(u_1, \dots, u_n)$ such that $n-i+1 \leq u_i \leq n$ for all i .

These sequences form an **interval** with $u_{\min} = (n, n-1, \dots, 1)$ and $u_{\max} = (n, n, \dots, n)$.

Proposition [NT 24]: the NT poset is a **lattice**.

Corollary. The ascent order on Dyck paths of size n defines a **lattice**.






$$v = (8, 8, 6, 6, 6, 6, 5, 5)$$

$$u = (8, 7, 6, 6, 6, 5, 3, 1)$$

The Nadeau-Tewari poset and the ascent order




Corollary. The ascent order on Dyck paths of size n defines a **lattice**.



The Nadeau-Tewari poset and the ascent order

Corollary. The ascent order on Dyck paths of size n defines a **lattice**.

Join(P,Q): the lowest Dyck path above P and Q whose descents are included in $\text{des}(P)$ and in $\text{des}(Q)$.






The Nadeau-Tewari poset and the ascent order

Corollary. The ascent order on Dyck paths of size n defines a **lattice**.

Join(P,Q): the lowest Dyck path above P and Q whose descents are included in $\text{des}(P)$ and in $\text{des}(Q)$.

Meet(P,Q): the highest Dyck path below P and Q whose descents contain those of P and Q.



III. The number of intervals


Interval $[P, Q] \sim (P, Q)$ with $P \leq Q$

Recursive construction of ascent intervals

Recursive construction of ascent intervals

Proposition. In the ascent poset, $P \preceq Q$ iff


- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



Recursive construction of ascent intervals

Proposition. In the ascent poset, $P \preceq Q$ iff


- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



Recursive construction of ascent intervals

Proposition. In the ascent poset, $P \preceq Q$ iff


- ◆ P lies below Q
- ◆ every descent of Q **is** (i.e. lies on the same diagonal as) a descent of P .



Corollary: if $[P, Q]$ is an interval, deleting the last peak of P and the last peak of Q gives a new interval.

Recursive construction of ascent intervals


Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...



Recursive construction of ascent intervals

Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...


- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$



Recursive construction of ascent intervals

Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...


- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$
- ◆ $a' \leq b'$



Recursive construction of ascent intervals

Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...


- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$
- ◆ $a' \leq b'$
- ◆ if $a' = a+1$ then $b' = b+1$.



Recursive construction of ascent intervals

Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...


- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$
- ◆ $a' \leq b'$
- ◆ if $a' = a+1$ then $b' = b+1$.




Recursive construction of ascent intervals

Conversely, starting from an interval $[P, Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...

- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$
- ◆ $a' \leq b'$
- ◆ if $a' = a+1$ then $b' = b+1$.



$$\begin{array}{c} \xrightarrow{i=a-1} \\ \xrightarrow{j=b-a} \end{array}$$




Recursive construction of ascent intervals

Conversely, starting from an interval $[P,Q]$ with final peaks at heights $a \leq b$, adding peaks in P and Q at heights a' and b' gives an interval iff...

- ◆ $1 \leq a' \leq a+1, \quad 1 \leq b' \leq b+1$
- ◆ $a' \leq b'$
- ◆ if $a' = a+1$ then $b' = b+1$.

Bijection intervals of size n
 \approx
quadrant walks of length $n-1$ starting from $(0,0)$
 \approx
quadrant walks of length n starting and ending at $(0,0)$




Recursive construction of ascent intervals

- Let $Q(t;x,y)=Q(x,y)$ be the GF of the associated quadrant walks:

$$Q(x,y) = \sum_w t^{|w|} x^{i(w)} y^{j(w)}.$$

Then the GF of ascent intervals is $G=tQ(1,1)=Q(0,0)-1$.

Bijection intervals of size n
 \approx
quadrant walks of length $n-1$ starting
from $(0,0)$
 \approx
quadrant walks of length n starting
and ending at $(0,0)$




Recursive construction of ascent intervals

- Let $Q(t;x,y)=Q(x,y)$ be the GF of the associated quadrant walks:

$$Q(x,y) = \sum_w t^{|w|} x^{i(w)} y^{j(w)}.$$

Then the GF of ascent intervals is $G=tQ(1,1)=Q(0,0)-1$.

- Step-by-step description of the walks:



Recursive construction of ascent intervals


- Let $Q(t;x,y)=Q(x,y)$ be the GF of the associated quadrant walks:

$$Q(x, y) = \sum_w t^{|w|} x^{i(w)} y^{j(w)}.$$

Then the GF of ascent intervals is $G=tQ(1,1)=Q(0,0)-1$.

- Step-by-step description of the walks:

$$Q(x, y) = 1 + txQ(x, y) + ty^2 \frac{xQ(x, y) - yQ(y, y)}{(x - y)(y - 1)} - t \frac{xQ(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$



A functional equation with two catalytic variables

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + txQ(x, y) + ty^2 \frac{xQ(x, y) - yQ(y, y)}{(x - y)(y - 1)} - t \frac{xQ(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

Walks in the quadrant: a much studied topic !

Very few algebraic cases.

A functional equation with two catalytic variables

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + txQ(x, y) + ty^2 \frac{xQ(x, y) - yQ(y, y)}{(x - y)(y - 1)} - t \frac{xQ(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

Walks in the quadrant: a much studied topic !

Very few algebraic cases.

Thm. Ascent intervals have an algebraic GF, namely

$$G = Z(1 - 2Z + 2Z^3), \quad \text{where} \quad Z = t(1 + Z)(1 + 2Z)^2.$$

Uses Tutte's invariants

[Bernardi, mbm, Raschel 17(a)]

A functional equation with two catalytic variables

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + txQ(x, y) + ty^2 \frac{xQ(x, y) - yQ(y, y)}{(x - y)(y - 1)} - t \frac{xQ(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

Walks in the quadrant: a much studied topic !

Very few algebraic cases.

Thm. Ascent intervals have an algebraic GF, namely

$$G = Z(1 - 2Z + 2Z^3), \quad \text{where} \quad Z = t(1 + Z)(1 + 2Z)^2.$$

Uses Tutte's invariants

[Bernardi, mbm, Raschel 17(a)]

Asymptotics:


$$g(n) \sim \kappa \mu^n n^{-7/2}, \quad \text{with} \quad \mu = \frac{11 + 5\sqrt{5}}{2}.$$

IV. m-Dyck paths, and mirrored m-Dyck paths

m-Dyck paths and mirrored m-Dyck paths

In an **m-Dyck path**, the length of all **ascents** is a multiple of m.

m=2





m-Dyck paths and mirrored m-Dyck paths

In an **m-Dyck path**, the length of all **ascents** is a multiple of m.

In a **mirrored m-Dyck path**, the length of all **descents** is a multiple of m.

m=2




m-Dyck paths and mirrored m-Dyck paths

In an **m-Dyck path**, the length of all **ascents** is a multiple of m .

In a **mirrored m-Dyck path**, the length of all **descents** is a multiple of m .


$m=2$



→ Study the order induced by the ascent order on m-Dyck paths and mirrored m-Dyck paths.


m-Dyck paths

m=2



m-Dyck paths

m=2



m-Dyck paths form an **interval** in the ascent lattice A_{mn} , with


$$\min = (U^m D^m)^n$$

$$\max = U^{mn} D^{mn}.$$

In particular, it is a **lattice**.

Mirrored m-Dyck paths

Mirrored m-Dyck paths only form a **join semi-lattice**.



What about intervals?

What about intervals?

Intervals in m-Dyck paths:

- ◆ Stanley lattice: **D-finite GF** (i.e., linear DE with pol. coeff's)

$$\frac{2(m+2)((m+1)n)!((m+1)(n+1))!}{n!(n+1)!(mn+2)!(m(n+2)+2)!}$$

- ◆ Tamari lattice: **algebraic GF** [mbm, Fusy, Préville-Ratelle 11]

$$\frac{m+1}{n(mn+1)} \binom{(m+1)^2n+m}{n-1}$$

Conj: Bergeron,
Préville-Ratelle

- ◆ Greedy Tamari lattice: **algebraic GF** [mbm, Chapoton 24]

$$\frac{(m+2)(m+1)^{n-1}}{(mn+1)(mn+2)} \binom{(m+1)n}{n}.$$

What about intervals?

Intervals in mirrored m-Dyck paths:

- ◆ Stanley lattice: **D-finite GF** (i.e., linear DE with pol coeffs)

$$\frac{2(m+2)((m+1)n)!((m+1)(n+1))!}{n!(n+1)!(mn+2)!(m(n+2)+2)!}$$

- ◆ Tamari lattice ?
- ◆ Greedy Tamari lattice ?

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + tx^m Q(x, y) + ty^2 \frac{x^m Q(x, y) - y^m Q(y, y)}{(x - y)(y - 1)} - t \frac{x^m Q(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + tx^m Q(x, y) + ty^2 \frac{x^m Q(x, y) - y^m Q(y, y)}{(x - y)(y - 1)} - t \frac{x^m Q(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + tx^m Q(x, y) + ty^2 \frac{x^m Q(x, y) - y^m Q(y, y)}{(x - y)(y - 1)} - t \frac{x^m Q(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + tx^m Q(x, y) + ty^2 \frac{x^m Q(x, y) - y^m Q(y, y)}{(x - y)(y - 1)} - t \frac{x^m Q(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2 + 4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2 + 4}}{m} \right)^m$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $tQ(1,1)$, where $Q(x,y)=Q(t;x,y)$ satisfies:

$$Q(x, y) = 1 + tx^m Q(x, y) + ty^2 \frac{x^m Q(x, y) - y^m Q(y, y)}{(x - y)(y - 1)} - t \frac{x^m Q(x, 1) - Q(1, 1)}{(x - 1)(y - 1)}.$$

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2 + 4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2 + 4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{m^2 + 2 - \sqrt{m^2 + 4}}{2m^2 + 6}}.$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2+4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2+4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{m^2 + 2 - \sqrt{m^2 + 4}}{2m^2 + 6}}.$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2+4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2+4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{m^2 + 2 - \sqrt{m^2 + 4}}{2m^2 + 6}}.$$

What about ascent intervals?

m-Dyck paths: delete the **final** large peak $U^m D^m \rightarrow$ quadrant walks.

For $m > 1$, the exponent α is **irrational**, and hence the GF of intervals **cannot be D-finite**.

[Bostan, Raschel, Salvy 14]

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$g_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2 + 4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2 + 4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{m^2 + 2 - \sqrt{m^2 + 4}}{2m^2 + 6}}.$$

What about ascent intervals?

Mirrored m-Dyck paths: delete the **first** large peak $U^m D^m \rightarrow$ quadrant walks.

What about ascent intervals?

Mirrored m-Dyck paths: delete the **first** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $t\bar{Q}(1,1)$, where $\bar{Q}(x,y)=\bar{Q}(t;x,y)$ satisfies:

$$\bar{Q}(x,y) = 1 + tx^m \frac{y\bar{Q}(x,y) - \bar{Q}(x,1)}{y-1} + ty^2 \frac{x^m \bar{Q}(x,y) - \bar{Q}(1,y)}{(x-1)(y-1)} - t \frac{x^m \bar{Q}(x,1) - \bar{Q}(1,1)}{(x-1)(y-1)}$$

What about ascent intervals?

Mirrored m-Dyck paths: delete the **first** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $t\bar{Q}(1,1)$, where $\bar{Q}(x,y)=\bar{Q}(t;x,y)$ satisfies:

$$\bar{Q}(x,y) = 1 + tx^m \frac{y\bar{Q}(x,y) - \bar{Q}(x,1)}{y-1} + ty^2 \frac{x^m \bar{Q}(x,y) - \bar{Q}(1,y)}{(x-1)(y-1)} - t \frac{x^m \bar{Q}(x,1) - \bar{Q}(1,1)}{(x-1)(y-1)}$$

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$\bar{g}_m(n) \sim \kappa \mu^n n^\alpha,$$

What about ascent intervals?

Mirrored m-Dyck paths: delete the **first** large peak $U^m D^m \rightarrow$ quadrant walks.

- The GF of ascent intervals is $t\bar{Q}(1,1)$, where $\bar{Q}(x,y)=\bar{Q}(t;x,y)$ satisfies:

$$\bar{Q}(x,y) = 1 + tx^m \frac{y\bar{Q}(x,y) - \bar{Q}(x,1)}{y-1} + ty^2 \frac{x^m \bar{Q}(x,y) - \bar{Q}(1,y)}{(x-1)(y-1)} - t \frac{x^m \bar{Q}(x,1) - \bar{Q}(1,1)}{(x-1)(y-1)}$$

- Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$\bar{g}_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2+4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2+4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{1 + 2m^2 - m\sqrt{1 + 4m^2}}{2(3m^2 + 1)}}.$$

What about ascent intervals?

Mirrored m-Dyck paths: delete the **first** large peak $U^m D^m \rightarrow$ quadrant walks.

For $m > 1$, the exponent α is **irrational**, and hence the GF of intervals **cannot be D-finite**.

[Bostan, Raschel, Salvy 14]

- **Asymptotics** (from random walk results) [Denisov & Wachtel 15]

$$\bar{g}_m(n) \sim \kappa \mu^n n^\alpha,$$

where

$$\mu = \frac{m\sqrt{m^2 + 4} + m^2 + 2}{2} \cdot \left(\frac{2 + \sqrt{m^2 + 4}}{m} \right)^m$$

and

$$\alpha = -1 - \pi / \arccos(c) \quad \text{with} \quad c = \sqrt{\frac{1 + 2m^2 - m\sqrt{1 + 4m^2}}{2(3m^2 + 1)}}.$$

V. Connection with the sylvester congruence

[Hivert, Novelli, Thibon 05]



In the OEIS...

Observation: for $m=1, 2, \dots, 5$, the sequence $\bar{g}_m(n)$ that counts intervals of mirrored m -Dyck paths appears in the OEIS.

In the OEIS...

Observation: for $m=1, 2, \dots, 5$, the sequence $\bar{g}_m(n)$ that counts intervals of mirrored m -Dyck paths appears in the OEIS.

- **$m=1$:** number of sylvester classes of 1-multiparking functions

Search: **seq:1,3,13,69,417,2759 id:243688**

Displaying 1-1 of 1 result found.

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#) Format: long | [short](#) | [data](#)

[A243688](#) Number of Sylvester classes of 1-multiparking functions of length n .

1, 3, 13, 69, 417, 2759

([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 1,2

COMMENTS See Novelli-Thibon (2014) for precise definition.

LINKS [Table of \$n, a\(n\)\$ for \$n=1..6\$.](#)

J.-C. Novelli, J.-Y. Thibon, [Hopf Algebras of \$m\$ -permutations, \$\(m+1\)\$ -ary trees, and \$m\$ -parking functions](#), arXiv preprint arXiv:1403.5962, 2014. See Fig. 26.

KEYWORD nonn,more

AUTHOR [N. J. A. Sloane](#), Jun 14 2014

STATUS approved

In the OEIS...

Observation: for $m=1, 2, \dots, 5$, the sequence $\bar{g}_m(n)$ that counts intervals of mirrored m -Dyck paths appears in the OEIS.

- **$m=1$:** number of sylvester classes of 1-multiparking functions
- **$m=2$:** number of sylvester classes of 2-multiparking functions

Search: **seq:1,5,40,407,4797 id:243671**

Displaying 1-1 of 1 result found.

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#) Format: long | [short](#) | [data](#)

[A243671](#) Number of Sylvester classes of 2-parking functions of length n .

1, 5, 40, 407, 4797

([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 1,2

COMMENTS See Novelli-Thibon (2014) for precise definition.

LINKS [Table of \$n, a\(n\)\$ for \$n=1..5\$.](#)

J.-C. Novelli, J.-Y. Thibon, [Hopf Algebras of \$m\$ -permutations, \$\(m+1\)\$ -ary trees, and \$m\$ -parking functions](#), preprint arXiv:1403.5962, 2014. See Fig. 21.

KEYWORD nonn,more

AUTHOR [N. J. A. Sloane](#), Jun 14 2014

STATUS approved

In the OEIS...

Observation: for $m=1, 2, \dots, 5$, the sequence $\bar{g}_m(n)$ that counts intervals of mirrored m -Dyck paths appears in the OEIS.

- **$m=1$:** number of sylvester classes of 1-multiparking functions
- **$m=2$:** number of sylvester classes of 2-multiparking functions
and so on.

Search: **seq:1,5,40,407,4797 id:243671**

Displaying 1-1 of 1 result found.

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#) Format: long | [short](#) | [data](#)

[A243671](#) Number of Sylvester classes of 2-parking functions of length n .

1, 5, 40, 407, 4797

([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 1,2

COMMENTS See Novelli-Thibon (2014) for precise definition.

LINKS [Table of \$n, a\(n\)\$ for \$n=1..5\$.](#)

J.-C. Novelli, J.-Y. Thibon, [Hopf Algebras of \$m\$ -permutations, \$\(m+1\)\$ -ary trees, and \$m\$ -parking functions](#), preprint arXiv:1403.5962, 2014. See Fig. 21.

KEYWORD nonn,more

AUTHOR [N. J. A. Sloane](#), Jun 14 2014

STATUS approved

In the OEIS...

Observation: for $m=1, 2, \dots, 5$, the sequence $\bar{g}_m(n)$ that counts intervals of mirrored m -Dyck paths appears in the OEIS.

- **$m=1$:** number of sylvester classes of 1-multiparking functions
 - **$m=2$:** number of sylvester classes of 2-multiparking functions
- and so on.

Link with the
NT-lattice

Search: **seq:1,5,40,407,4797 id:243671**

Displaying 1-1 of 1 result found.

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#) Format: long | [short](#) | [data](#)

[A243671](#) Number of Sylvester classes of 2-parking functions of length n .

1, 5, 40, 407, 4797

([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 1,2

COMMENTS See Novelli-Thibon (2014) for precise definition.

LINKS [Table of \$n, a\(n\)\$ for \$n=1..5\$.](#)

J.-C. Novelli, J.-Y. Thibon, [Hopf Algebras of \$m\$ -permutations, \$\(m+1\)\$ -ary trees, and \$m\$ -parking functions](#), preprint arXiv:1403.5962, 2014. See Fig. 21.

KEYWORD nonn,more

AUTHOR [N. J. A. Sloane](#), Jun 14 2014

STATUS approved

The sylvester congruence

- Defined on words on the alphabet \mathbb{Z}
- Generated by commutation relations:

$$ac \cdots b \equiv ca \cdots b, \quad a \leq b < c.$$

- Class representatives: words avoiding **aba** and **acb** (as subwords), called **sylvester words**.

The sylvester congruence

- Defined on words on the alphabet \mathbb{Z}
- Generated by commutation relations:

$$ac \cdots b \equiv ca \cdots b, \quad a \leq b < c.$$

- Class representatives: words avoiding **aba** and **acb** (as subwords), called **sylvester words**.

Example:

2 4 **3 5 3** \rightarrow 2 4 **5 3 3**

The sylvester congruence

- Defined on words on the alphabet \mathbb{Z}
- Generated by commutation relations:

$$ac \cdots b \equiv ca \cdots b, \quad a \leq b < c.$$

- Class representatives: words avoiding **aba** and **acb** (as subwords), called **syvester words**.

Example:

$$2\ 4\ 3\ 5\ 3 \rightarrow 2\ 4\ 5\ 3\ 3$$

$$2\ 4\ 5\ 3\ 3 \rightarrow 4\ 2\ 5\ 3\ 3$$

The sylvester congruence

- Defined on words on the alphabet \mathbb{Z}
- Generated by commutation relations:

$$ac \cdots b \equiv ca \cdots b, \quad a \leq b < c.$$

- Class representatives: words avoiding **aba** and **acb** (as subwords), called **sylvester words**.

Example:

$$2\ 4\ 3\ 5\ 3 \rightarrow 2\ 4\ 5\ 3\ 3$$

$$2\ 4\ 5\ 3\ 3 \rightarrow 4\ 2\ 5\ 3\ 3$$

$$4\ 2\ 5\ 3\ 3 \rightarrow 4\ 5\ 2\ 3\ 3, \text{ sylvester word}$$

The Nadeau-Tewari lattice


[2024]

Def. Let $u=(u_1, \dots, u_n)$ and $v=(v_1, \dots, v_n)$ be two nonincreasing sequences of integers. Then $u \leq v$ for the NT order if

- u lies below v ($u_i \leq v_i$)
- every descent of v is a descent of u .

$$v = (10, 10, 10, 10, 9, 4)$$

$$u = (10, 10, 8, 8, 7, 3)$$



From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 5$.

From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$.

Let $w_1 = \text{Ninc}(w) = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ be its nonincreasing reordering.

From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$.

Let $w_1 = \text{Ninc}(w) = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ be its nonincreasing reordering.

Let $w_2 = \text{LRMin}(w) = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$ be the largest nonincreasing word that is smaller than w , componentwise.


From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$.

Let $w_1 = \text{Ninc}(w) = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ be its nonincreasing reordering.

Let $w_2 = \text{LRMin}(w) = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$ be the largest nonincreasing word that is smaller than w , componentwise.

- Write w_1 and w_2 vertically as follows:




From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$.

Let $w_1 = \text{Ninc}(w) = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ be its nonincreasing reordering.

Let $w_2 = \text{LRMin}(w) = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$ be the largest nonincreasing word that is smaller than w , componentwise.

- Write w_1 and w_2 vertically as follows:
- Complete with $n=6$ horizontal steps to form two ES paths.




From sylvester words to Nadeau-Tewari intervals

Example. Fix $n=6$ and a sylvester word w on the alphabet $\{1, 2, \dots, n\}$, containing the letter 1, say $w = 3 2 2 2 2 5 1 1 1 5$.

Let $w_1 = \text{Ninc}(w) = 5 5 3 2 2 2 2 1 1 1$ be its nonincreasing reordering.

Let $w_2 = \text{LRMin}(w) = 3 2 2 2 2 2 1 1 1 1$ be the largest nonincreasing word that is smaller than w , componentwise.

- Write w_1 and w_2 vertically as follows:
- Complete with $n=6$ horizontal steps to form two ES paths.
- The horizontal words $u = 10 10 8 8 7 3$ and $v = 10 10 10 10 9 4$, of length $n=6$, form an interval in the Nadeau-Tewari lattice.




From sylvester words to Nadeau-Tewari intervals

Proposition. For any n , this construction gives a bijection between:

- ◆ sylvester words w on the alphabet $\{1, 2, \dots, n\}$ containing the letter 1, and
- ◆ intervals $[u,v]$ in the NT lattice, such that u and v have length n , **positive entries** and the **same first letter**.

Example

For $n=6$ and $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$, we have $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$.



From sylvester words to Nadeau-Tewari intervals


Proposition. For any n , this construction gives a bijection between:

- ◆ sylvester words w on the alphabet $\{1, 2, \dots, n\}$ containing the letter 1, and
- ◆ intervals $[u,v]$ in the NT lattice, such that u and v have length n , **positive entries** and the **same first letter**.

Example

For $n=6$ and $w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$, we have $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$.


Conversely?



From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:



From Sylvester words to Nadeau-Tewari intervals


Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.



From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = _ _ _ _ _ _ _ _ _ _ _$

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ _ _ _ _ 1\ _ _ _$

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ _ _ _ _ 1\ _ _ _$

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ _ _ _ _ 1\ _ _ _ _$

non-dec.

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ _ _ _$

non-dec.

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ _ _ _$

non-dec.

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Conversely?

- Starting from $u = 10\ 10\ 8\ 8\ 7\ 3$ and $v = 10\ 10\ 10\ 10\ 9\ 4$, recover the vertical words $w_1 = 5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1$ and $w_2 = 3\ 2\ 2\ 2\ 2\ 2\ 1\ 1\ 1\ 1$:

same length (10),
contain 1,

w_2 is less than w_1 (componentwise)

$\text{alph}(w_2) \subset \text{alph}(w_1)$.

- Recover w from $w_1 = \text{NInc}(w)$
and $w_2 = \text{LRMin}(w)$?

$\text{lett}(w) = \{5\ 5\ 3\ 2\ 2\ 2\ 2\ 1\ 1\ 1\}$

$w = 3\ 2\ 2\ 2\ 2\ 5\ 1\ 1\ 1\ 5$

non-dec.

avoids aba and acb

From sylvester words to Nadeau-Tewari intervals

Proposition. For any n , this construction gives a bijection between:

- ◆ sylvester words w on the alphabet $\{1, 2, \dots, n\}$ containing the letter 1, and
- ◆ intervals $[u,v]$ in the NT lattice, such that u and v have length n , **positive entries** and the **same first letter**.

From sylvester words to Nadeau-Tewari intervals

Proposition. For any n , this construction gives a bijection between:

- ♦ sylvester words w on the alphabet $\{1, 2, \dots, n\}$ containing the letter 1 , and
- ♦ intervals $[u, v]$ in the NT lattice, such that u and v have length n , **positive entries** and the **same first letter**.

Specializations: bijections between

- ♦ positive sylvester words w of length mn such that $N\text{Inc}(w) \leq n^m (n-1)^m \dots 2^m 1^m$ and **ascent intervals of m -Dyck paths** of length mn
- ♦ positive sylvester words w of length n such that $N\text{Inc}(w) \leq ((n-1)m+1) \dots (2m+1) (m+1) 1$ and **ascent intervals of mirrored m -Dyck paths** of length mn .

VI. Final remarks

Final remarks

- **Combinatorial proof** for the number/GF of ascent intervals? ($m=1$)

$$(n + 4) (2n + 7) g(n + 2) = 2 (11n^2 + 44n + 42) g(n + 1) + n (2n + 1) g(n)$$

Final remarks

- **Combinatorial proof** for the number/GF of ascent intervals? ($m=1$)

$$(n + 4) (2n + 7) g(n + 2) = 2 (11n^2 + 44n + 42) g(n + 1) + n (2n + 1) g(n)$$


- **A symmetric joint distribution** on ascent intervals $[P, Q]$ ($m=1$):

$a(P)$ = length of the first ascent of P

$r(P, Q)$ = number of ascents of P before the first descent of Q

Example. $a(P)=1$, $r(P, Q)=2$

(Non-recursive) **bijection?**



Final remarks

- **Combinatorial proof** for the number/GF of ascent intervals? ($m=1$)

$$(n + 4) (2n + 7) g(n + 2) = 2 (11n^2 + 44n + 42) g(n + 1) + n (2n + 1) g(n)$$


- **A symmetric joint distribution** on ascent intervals $[P, Q]$ ($m=1$):

$a(P)$ = length of the first ascent of P

$r(P, Q)$ = number of ascents of P before the first descent of Q

Example. $a(P)=1$, $r(P, Q)=2$

(Non-recursive) **bijection?**



- Study **mirrored m -Dyck paths** in other Dyck lattices: intervals?

Final remarks

- **Combinatorial proof** for the number/GF of ascent intervals? ($m=1$)

$$(n + 4) (2n + 7) g(n + 2) = 2 (11n^2 + 44n + 42) g(n + 1) + n (2n + 1) g(n)$$


- **A symmetric joint distribution** on ascent intervals $[P, Q]$ ($m=1$):

$a(P)$ = length of the first ascent of P

$r(P, Q)$ = number of ascents of P before the first descent of Q

Example. $a(P)=1$, $r(P, Q)=2$

(Non-recursive) **bijection?**



- Study **mirrored m-Dyck paths** in other Dyck lattices: intervals?
- Poset properties? (shellability, geometric realizations...)

Final remarks

- **Combinatorial proof** for the number/GF of ascent intervals? ($m=1$)

$$(n+4)(2n+7)g(n+2) = 2(11n^2 + 44n + 42)g(n+1) + n(2n+1)g(n)$$

- **A symmetric joint distribution** on ascent intervals $[P, Q]$ ($m=1$):

$a(P)$ = length of the first ascent of P

$r(P, Q)$ = number of ascents of P before the first descent of Q

Example. $a(P)=1$, $r(P, Q)=2$


(Non-recursive) **bijection**?

Thanks for your
attention

- Study **mirrored m -Dyck paths** in other Dyck lattices: intervals?
- Poset properties? (shellability, geometric realizations...)

m-Dyck paths and mirrored m-Dyck paths

$m=2$



m-Dyck paths and mirrored m-Dyck paths

$m=2$

